



TRILYTE PREP – AM Procedure

**** Nothing by mouth within 3 hours of your procedure or it may be canceled. ****

A Colonoscopy is an examination of the colon (large bowel) with a flexible lighted tube about the size of your ring finger which transmits a live color image onto a television screen. A good bowel prep is essential for an accurate and successful procedure. These instructions will help maximize the ease, safety, and success of your preparation.

You must have a responsible adult accompany you for the appointment. We ask that you also have a responsible adult at home with you following the procedure. Please be aware you **will not** be permitted to drive home by yourself or leave unaccompanied in a taxi or Uber.

Prior to your Colonoscopy you will need to purchase the following items:

- **Trilyte bowel prep. This has been called in to your pharmacy. Please pick it up at your convenience.**
- Clear liquids of choice: fruit juices without pulp i.e. apple, white grape, water, clear broth bouillon, coffee without cream, soft drinks/soda, jello, popsicles – nothing red/purple

A word about bowel preps:

- NO bowel prep is fun.
- The purpose of the bowel prep is to clean out the colon to allow clear visualization for the exam.
- A bowel prep is the only way to effectively clean out the bowel (if there was another way, we would use it).
- It is a short-term discomfort for a long-term cause.
- Your bowel movements may be frequent, explosive, and liquid (this is to be expected). Respond immediately to the urge.
- Use soft toilet paper and zinc oxide or A&D ointment for skin irritation.

Preparing for Your Colonoscopy

Two days before your procedure:

- Two days before your procedure start clear liquids at 6:00am. Do not eat solid foods.

One day before your procedure:

- Start clear liquids at 6:00 am and stay well hydrated throughout the day. Do not eat food. Prepare the Trilyte as directed and place it in the refrigerator. Some of these products come with a Flavor Pack. If you wish to use it, add the contents of the Flavor Pack into the bottle before you add the water. (Using the flavor pack is optional)
- The evening prior to the procedure, drink 2 liters (half of the jug) by drinking 8 oz. every 10 minutes. If you start at 6:00 pm, then expect to finish drinking 2 liters by 8 pm. Stay near a bathroom and remain hydrated with clear liquids. Medications (except blood thinners) may be taken and teeth may be brushed.

The day of your procedure:

- Five (5) hours before your procedure, drink the remaining 2 liters (8 oz. every 10 min.) Remain hydrated on clear liquids until 3 hours prior to the exam, at which point nothing else should be ingested (taken by mouth). Plan ahead by finishing the Trilyte at home, allowing for bowel movements, and adjusting for travel time.

Note: For patients with an artificial heart valve, pacemaker, an internal defibrillator, or those taking Coumadin, Wafarin, or Plavix, please consult with your physician regarding specific instructions.

Frequently Asked Questions

What is a clear stool?

A clear stool can have a slight tint of yellow or brown. It will be completely transparent and will not contain any solid matter. At the completion of the prep if it is brown or solid, please contact scheduling to discuss the situation.

I am not having bowel movements, what should I do?

Bowel movements can take up to 5-6 hours to start after beginning the prep. Be patient, continue to drink liquids, and proceed with the second dose of the bowel prep as directed. If you are still not seeing results, you will need to reach your physician's office after 7:30 AM for further instructions.

The prep is making me nauseated, what should I do?

If you develop nausea or vomiting, slow down the rate at which you drink the solution. Please attempt to drink all of the laxative solution even if it takes you longer. If vomiting persists, or you are not able to finish the preparation, stop the preparation and call your physician's office for further instructions.

What are some high fiber foods I should avoid?

Raw fruits and vegetables are typically high in fiber, as well as nuts, seeds, whole grain breads and beans and lentils.

What are some good options for low fiber foods?

Choose white bread and white rice for lower fiber options, as well as pastas made with white flour. Chicken, fish, dairy and eggs are also low in fiber and good choices for foods 2-3 days before you begin your prep.

If I eat popcorn or seeds 3 days before my procedure do I need to reschedule?

You will not need to reschedule your procedure, however the seeds or nuts may cause a difficulty in screening and require a need for rescreening. If you have eaten a large amount of seeds or nuts, you may want to contact the nurse or physician.

What is clear liquid?

Clear liquids you may have: Clear broth, apple juice, white grape juice, colas, coffee, tea, Jello, popsicles, water or any other clear liquid that if you hold it up to light you can see through it. Avoid the colors red and purple. Coffee creamer and milk or milk products are not considered a clear liquid.

Can I drink ALCOHOL on the liquid diet?

Alcohol is not allowed as part of the liquid diet.

Can I continue to be on the liquid diet after I begin consuming the laxatives?

Yes, you may continue the liquid diet until you are directed to discontinue anything by mouth, which is typically 3 hours prior to the procedure.

Why do I have to wake up at so early for the 2nd dose, can't I take it all the night before?

A split prep has proven to be the most effective for a successful colonoscopy. It is essential that you follow the directions provided with your prep medications.

If I weigh under 100 pounds do I need to take all of the prep?

The liquid amount is not weight dependent. It is important to finish the prep for a successful colonoscopy.

Note: If your insurance requires a pre-authorization, please keep in mind that obtaining this authorization does not guarantee payment. Contact your insurance for more information prior to your procedure.